


## Teen Pregnancy Rate, Jefferson County and Washington State, 1992-94 through 2009-11

Teen pregnancy rate is the number of live births and abortions to female residents age 15-17 per 1,000 female residents age 15-17 in a given place in a given time period.

Time Period	Jefferson County			95% confidence interval		WA Teen Pregnancy Rate per 1,000 females age 15-17
	pregnancies	population	Teen Pregnancy Rate per 1,000 females age 15-17	lower	upper	
1992-94	50	1166	<b>42.9</b>	31.9	56.6	<b>54.8</b>
1995-97*	46	1384	<b>33.2</b>	24.4	44.4	<b>46.2</b>
1998-00	59	1497	<b>39.4</b>	30.0	50.9	<b>38.4</b>
2001-03	48	1490	<b>32.2</b>	23.8	42.7	<b>30.8</b>
2004-06*	27	1453	<b>18.6</b>	12.3	27.0	<b>27.9</b>
2007-09*	20	1396	<b>14.3</b>	8.8	22.1	<b>26.7</b>
2009-11*	14	1271	<b>11.0</b>	6.0	18.4	<b>21.5</b>



\*WA rate statistically higher than Jefferson rate

## Teen Pregnancies, Jefferson County, 2000 through 2011

Teen pregnancies are the number of live births and abortions to Jefferson County female residents age 15-17

year	pregnancies	population	ratio of pregnant females age 15-17 to population of females age 15-17	
			Jefferson County	Washington State
2000	21	505	1 in 25	1 in 25
2001	20	501	1 in 25	1 in 30
2002	11	498	1 in 50	1 in 30
2003	17	491	1 in 30	1 in 35
2004	10	482	1 in 50	1 in 35
2005	10	487	1 in 50	1 in 35
2006	7	484	1 in 80	1 in 35
2007	5	479	1 in 100	1 in 35
2008	10	470	1 in 50	1 in 35
2009	5	447	1 in 90	1 in 40
2010	5	430	1 in 85	1 in 50
2011	4	394	1 in 100	1 in 60

**About 1 in 100 Jefferson females age 15-17 was pregnant in 2011**

Data Sources:

Vital Statistics Databases: Jefferson County Births and Abortions 1980-1989

WA State Dept of Health, Center for Health Statistics, Community Health Assessment Tool (CHAT) 1990-2011

Population estimates, 1983-1999 WA Office of Financial Management, 2000-2010, CHAT


2011 Population Estimate, Office of Financial Management

## Chlamydia & Gonorrhea Diagnosis Rates, Teens age 15-19, Jefferson County and Washington State

Year	Jefferson County FEMALES, age 15-19			95% confidence interval		WA Chlamydia rate per 1,000 females age 15-19
	chlamydia	population	Chlamydia rate per 1,000 females age 15-19	lower	upper	
2000	12	736	16.3	8.4	28.4	20.3
2001	5	733	6.8	2.2	15.7	20.5
2002	12	731	16.4	8.5	28.6	21.2
2000-02*	29	2200	13.2	8.8	18.9	20.7
2003	14	724	19.3	10.6	32.4	23.2
2004	13	717	18.1	9.7	30.9	23.8
2005	17	723	23.5	13.7	37.6	22.5
2003-05	44	2164	20.3	14.8	27.3	23.2
2006	8	719	11.1	4.8	21.8	21.6
2007	6	712	8.4	3.1	18.1	21.2
2008	19	707	26.9	16.2	41.9	23.7
2006-08*	33	2138	15.4	10.6	21.7	22.2
2009	17	684	24.9	14.5	39.7	24.4
2010	17	658	25.8	15.1	41.3	23.4
2011	10	659	15.2	7.3	27.8	24.8
2009-11	44	2001	22.0	16.0	29.5	24.2

Jefferson female age 15-19 gonorrhea diagnoses	WA Gonorrhea rate per 1,000 females age 15-19
0	1.8
0	2.2
0	1.9
0	2.0
0	1.8
0	2.1
0	2.4
0	2.1
0	2.8
0	2.5
0	2.1
0	2.4
0	1.2
0	1.4
1	1.3
1	1.3

**2 in 100 Jefferson female residents age 15-19 was diagnosed with Chlamydia in 2009-2011**



\*WA female Chlamydia rate statistically higher than Jefferson

Data sources: Washington State Dept. of Health, STD Data, Accessed in: Community Health Assessment Tool (CHAT). All case counts and denominators updated March 2013  
2011 denominators from OFM until CHAT denominators become available

## Chlamydia & Gonorrhea Diagnosis Rates, Teens age 15-19, Jefferson County and Washington State

Year	Jefferson County MALES, age 15-19			95% confidence interval		WA Chlamydia rate per 1,000 males age 15-19	Jefferson male age 15-19 gonorrhea diagnoses	WA gonorrhea rate per 1,000 males age 15-19
	chlamydia population	Chlamydia Rate per 1,000 males age 15-19		lower	upper			
2000	1	801				3.8	0	0.7
2001	0	796				3.3	0	0.8
2002	0	786				3.8	0	0.8
2000-02*	1	2383	n<5			3.6	0	0.8
2003	3	775				4.0	1	0.6
2004	2	765				4.3	1	0.7
2005	2	765				4.1	0	0.9
2003-05*	7	2305	3.0	1.2	6.2	4.1	2	0.7
2006	0	763				3.6	0	1.0
2007	1	757				3.9	0	1.0
2008	2	746				4.3	0	0.7
2006-08*	3	2266	n<5			3.9	0	0.9
2009	2	722				4.5	0	0.5
2010	4	696				4.4	0	0.5
2011	0	670				4.4	0	0.5
2009-11*	6	2088	2.9	1.1	6.2	4.4	0	0.5

<p><b>3 in 1,000 Jefferson males age 15-19 was diagnosed with Chlamydia in 2009-11</b></p>	<p>*WA male Chlamydia rate statistically higher than Jefferson.</p>
--	---

Data sources: Washington State Dept. of Health, STD Data, Accessed in: Community Health Assessment Tool (CHAT). All case counts and denominators updated March 2013  
2011 denominators from OFM until CHAT denominators become available

**General Health, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012**

8th grade	2004				2006				2008				2010				2012								
	response rate: 72%			95% confidence interval	51%			95% confidence interval	88%			95% confidence interval	70%			95% confidence interval	82%			95% confidence interval					
	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper					
physical or check-up in the past 12 months	51	72	71%	53%	93%	47	66	71%	52%	95%	43	86	50%	36%	67%	47	71	66%	49%	88%	not asked				
moderate exercise, 5 or more days in past week	34	94	36%	25%	51%	35	76	46%	32%	64%	35	89	39%	27%	55%	not asked					not asked				
vigorous exercise, 3 or more days in past week	71	94	76%	59%	95%	62	76	82%	63%	105%	73	90	81%	64%	100%	not asked					not asked				
physically active 60+ minutes/day 7 days/week	not asked					22	75	29%	18%	44%	22	88	25%	16%	38%	30	74	41%	27%	58%	25	91	28%	18%	41%
never/seldom/sometimes enjoyed school	71	201	35%	28%	45%	58	149	39%	30%	50%	108	179	60%	50%	73%	78	151	52%	41%	65%	92	180	51%	41%	63%
did not feel safe at school	47	203	23%	17%	31%	32	149	21%	15%	30%	46	187	25%	18%	33%	25	157	16%	10%	24%	25.8	182	14%	9%	21%
family cut meal size or skipped meals in past year	12	81	15%	8%	26%	10	38	26%	13%	48%	12	73	16%	9%	29%	8	62	13%	6%	25%	15	88	17%	10%	28%
overweight or obese	20	84	24%	15%	37%	16	67	24%	14%	39%	12	77	16%	8%	27%	13	61	21%	11%	36%	24.1	83	29%	19%	43%
had 1 or more sodas yesterday	56	96	58%	44%	76%	30	76	39%	27%	56%	38	90	42%	30%	58%	22	74	30%	19%	45%	37	91	41%	29%	56%
have 5 or more fruits/vegetables a day	21	94	22%	14%	34%	29	70	41%	28%	60%	26	78	33%	22%	49%	not asked					29	89	33%	22%	47%
most of the time or always have dinner with family	63	96	66%	50%	84%	52	76	68%	51%	90%	65	90	72%	56%	92%	55	75	73%	55%	96%	62	92	67%	52%	86%
ate breakfast today	not asked					48	76	63%	47%	84%	63	90	70%	54%	90%	61	75	81%	62%	100%	54	91	59%	45%	77%
in at least one physical fight in past year	63	204	31%	24%	40%	55	148	37%	28%	48%	72	187	39%	30%	49%	57	158	36%	27%	47%	65	179	36%	28%	46%
bullied at least once in the past 30 days	71	202	35%	27%	44%	45	144	31%	23%	42%	52	177	29%	22%	38%	59	154	38%	29%	49%	76	181	42%	33%	53%
bullied on computer or cell phone in past 30 days	not asked					6	74	8%	3%	17%	10	94	11%	5%	19%	14	70	20%	11%	33%	11	83	13%	7%	24%
taught about HIV infection at school last year	not asked					61	75	81%	62%	100%	67	84	80%	62%	100%	57	67	85%	64%	100%	74	89	83%	65%	100%

Response Rates:

70% or greater participation: results probably representative

40-69% participation: results may be representative

Less than 40% participation: results likely not representative

**General Health, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012**

10th grade	2004				2006				2008				2010				2012								
	response rate: 61%			95% confidence interval	49%			95% confidence interval	53%			95% confidence interval	58%			95% confidence interval	53%			95% confidence interval					
	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper					
physical or check-up in the past 12 months	59	81	73%	55%	94%	36	68	53%	37%	73%	37	68	54%	38%	75%	36	69	52%	37%	72%	not asked				
moderate exercise, 5 or more days in past week	29	90	32%	22%	46%	29	76	38%	26%	55%	27	67	40%	27%	59%	not asked					not asked				
vigorous exercise, 3 or more days in past week	59	90	66%	50%	85%	55	77	71%	54%	93%	45	66	68%	50%	91%	not asked					not asked				
physically active 60+ minutes/day 7 days/week	not asked					16	78	21%	12%	33%	18	66	27%	16%	43%	15	69	22%	12%	36%	13	54	24%	13%	41%
never/seldom/sometimes enjoyed school	59	176	34%	26%	43%	46	138	33%	24%	44%	82	142	58%	46%	72%	98	156	63%	51%	77%	67.9	111	61%	48%	78%
did not feel safe at school	38	182	21%	15%	29%	24	150	16%	10%	24%	15	141	11%	6%	18%	19	150	13%	8%	20%	22	113	20%	12%	30%
family cut meal size or skiped meals in past year	14	74	19%	10%	32%	10	30	33%	16%	61%	10	64	16%	7%	29%	20	67	30%	18%	46%	9.02	54	17%	8%	31%
overweight or obese	23	89	26%	16%	39%	20	76	26%	16%	41%	16	66	24%	14%	39%	12	65	19%	10%	32%	7.99	49	16%	7%	32%
had 1 or more sodas yesterday	39	90	43%	31%	59%	33	78	42%	29%	59%	22	67	33%	21%	50%	19	69	28%	17%	43%	19	54	35%	21%	55%
have 5 or more fruits/vegetables a day	32	89	36%	25%	51%	14	53	26%	14%	44%	19	65	29%	18%	46%	not asked					18	53	34%	20%	54%
most of the time or always have dinner with family	46	89	52%	38%	69%	46	77	60%	44%	80%	50	67	75%	55%	98%	44	69	64%	46%	86%	33	54	61%	42%	86%
ate breakfast today	not asked					54	78	69%	52%	90%	40	67	60%	43%	81%	39	69	57%	40%	77%	34	54	63%	44%	88%
in at least one physical fight in past year	62	184	34%	26%	43%	53	150	35%	26%	46%	44	142	31%	23%	42%	32	150	21%	15%	30%	29.9	113	27%	18%	38%
bullied at least once in the past 30 days	46	178	26%	19%	34%	37	133	28%	20%	38%	35	137	26%	18%	35%	32	144	22%	15%	31%	38	113	34%	24%	46%
bullied on computer or cell phone in past 30 days	not asked					7	72	10%	4%	20%	8	68	12%	5%	23%	8	69	12%	5%	23%	11	54	20%	10%	36%
taught about HIV infection at school last year	not asked					52	67	78%	58%	100%	54	68	79%	60%	100%	61	68	90%	69%	100%	45	54	83%	61%	100%

Data Sources: Washington State Department of Health, Healthy Youth Survey

Response Rates:

70% or greater participation: results probably representative

40-69% participation: results may be representative

Less than 40% participation: results likely not representative

## Mental Health, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012

8th grade	2004			95% confidence interval		2006	2008			95% confidence interval		2010	2012			95% confidence interval									
	response rate: 72%	#	total	%	lower		upper	51%	#	total	%		lower	upper	70%	#	total	%	lower	upper					
felt so sad or hopeless for at least 2 weeks, stopped usual activities	66	197	34%	26%	43%	38	143	27%	19%	36%	34	173	20%	14%	27%	47	150	31%	23%	42%	45	180	25%	18%	33%
seriously considered suicide in past year	18	89	20%	12%	32%	8	76	11%	5%	21%	25	173	14%	9%	21%	28	148	19%	13%	27%	27.1	178	15%	10%	22%
made plan for suicide in past year	13	98	13%	7%	23%	8	76	11%	5%	21%	8	92	9%	4%	17%	6	75	8%	3%	17%	14	92	15%	8%	25%
when feeling sad or hopeless, have adults to go to for help	45	76	59%	43%	79%	38	55	69%	49%	95%	39	64	61%	43%	83%	32	57	56%	38%	79%	51	69	74%	55%	97%
very/somewhat likely to seek help if depressed/suicidal	23	46	50%	32%	75%	11	30	37%	18%	65%	27	43	63%	41%	91%	19	33	58%	35%	90%	23	43.03	53%	34%	80%
very/somewhat likely to seek help for friend if depressed/suicidal	70	91	77%	60%	97%	61	74	82%	63%	106%	64	85	75%	58%	96%	51	68	75%	56%	99%	73.9	87	85%	67%	107%

10th grade	2004			95% confidence interval		2006	2008			95% confidence interval		2010	2012			95% confidence interval									
	response rate: 61%	#	total	%	lower		upper	49%	#	total	%		lower	upper	53%	#	total	%	lower	upper					
felt so sad or hopeless for at least 2 weeks, stopped usual activities	63	176	36%	28%	46%	36	120	30%	21%	42%	34	137	25%	17%	35%	44	147	30%	22%	40%	39	113	35%	25%	47%
seriously considered suicide in past year	20	96	21%	13%	32%	15	80	19%	11%	31%	19	138	14%	8%	22%	28	146	19%	13%	28%	27	113	24%	16%	35%
made plan for suicide in past year	14	96	15%	8%	24%	12	80	15%	8%	26%	6	68	9%	3%	19%	8	69	12%	5%	23%	8.01	56	14%	6%	28%
when feeling sad or hopeless, have adults to go to for help	57	79	72%	55%	94%	34	66	52%	36%	72%	39	55	71%	50%	97%	34	58	59%	41%	82%	31	56	55%	38%	79%
very/somewhat likely to seek help if depressed/suicidal	26	53	49%	32%	72%	26	46	57%	37%	83%	30	44	68%	46%	97%	25	46.02	54%	35%	80%	19	34.99	54%	33%	85%
very/somewhat likely to seek help for friend if depressed/suicidal	66	88	75%	58%	95%	60	73	82%	63%	100%	56	68	82%	62%	100%	60	68	88%	67%	100%	46	54	85%	62%	100%

Data Sources: Washington State Department of Health, Healthy Youth Survey

### Response Rates:

70% or greater participation: results probably representative

40-69% participation: results may be representative

Less than 40% participation: results likely not representative

## Oral Health, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012

8th grade	2004				2006				2008				2010				2012								
	response rate: 72%			95% confidence interval		51%			95% confidence interval		88%			95% confidence interval		70%			95% confidence interval		82%			95% confidence interval	
	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper
dental check-up past 12 months	69	85	81%	63%	100%	58	68	85%	65%	100%	60	82	73%	56%	94%	52	71	73%	55%	96%	66	90	73%	57%	93%

10th grade	2004				2006				2008				2010				2012								
	response rate: 61%			95% confidence interval		49%			95% confidence interval		53%			95% confidence interval		58%			95% confidence interval		53%			95% confidence interval	
	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper
dental check-up past 12 months	61	85	72%	55%	92%	51	69	74%	55%	97%	43	68	63%	46%	85%	48	69	70%	51%	92%	39	54	72%	51%	99%

Data Sources: Washington State Department of Health, Healthy Youth Survey

### Response Rates:

70% or greater participation: results probably representative

40-69% participation: results may be representative

Less than 40% participation: results likely not representative

## Chronic Disease, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012

8th grade	2004				2006				2008				2010				2012								
	response rate: 72%			95% confidence interval		51%			95% confidence interval		88%			95% confidence interval		70%			95% confidence interval		82%			95% confidence interval	
	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper
ever told by doctor have asthma	20	90	22%	14%	34%	8	69	12%	5%	23%	12	88	14%	7%	24%	12	71	17%	9%	29%	17	92	19%	11%	30%
ever told by doctor have diabetes	6	72	8%	3%	18%	3	69	4%	1%	12%	4	76	5%	1%	13%	3	63	5%	1%	13%	not asked				

10th grade	2004				2006				2008				2010				2012								
	response rate: 61%			95% confidence interval		49%			95% confidence interval		53%			95% confidence interval		58%			95% confidence interval						
	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper	#	total	%	lower	upper
ever told by doctor have asthma	11	85	13%	6%	23%	14	71	20%	11%	33%	16	68	24%	13%	38%	15	69	22%	12%	36%	13	54	24%	13%	41%
ever told by doctor have diabetes	5	70	7%	2%	16%	3	40	8%	2%	21%	4	66	6%	2%	15%	1	69	2%	0%	7%	not asked				

Data Sources: Washington State Department of Health, Healthy Youth Survey

### Response Rates:

70% or greater participation: results probably representative

40-69% participation: results may be representative

Less than 40% participation: results likely not representative



# Substance Use, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012

8th grade	2004			95% confidence interval		2006			95% confidence interval		2008			95% confidence interval		2010			95% confidence interval		2012			95% confidence interval						
	response rate: 72%	#	total	%	lower	upper	51%	#	total	%	lower	upper	88%	#	total	%	lower	upper	70%	#	total	%	lower	upper	82%	#	total	%	lower	upper
ever had more than a sip or two of alcohol		108	200	54%	44%	65%		63	148	43%	33%	54%		78	180	43%	34%	54%		63	154	41%	31%	52%		77	180	43%	34%	54%
binge drank at least once in past 2 weeks		41	202	20%	15%	28%		6	72	8%	3%	18%		24	179	13%	9%	20%		15	158	9%	5%	16%		18	180	10%	6%	16%
drunk at school in past year		26	202	13%	8%	19%		14	148	9%	5%	16%		19	181	10%	6%	16%		15	156	10%	5%	16%		13	179	7%	4%	12%
drank alcohol in past 30 days		58	202	29%	22%	37%		26	149	17%	11%	26%		36	183	20%	14%	27%		26	159	16%	11%	24%		22	180	12%	8%	18%
ever smoked marijuana		48	202	24%	18%	32%		30	148	20%	14%	29%		29	183	16%	11%	23%		30	156	19%	13%	27%		30	180	17%	11%	24%
smoked marijuana in past 30 days		38	203	19%	13%	26%		15	149	10%	6%	17%		20	183	11%	7%	17%		21	159	13%	8%	20%		22	180	12%	8%	18%
ever smoked a cigarette (even a puff)		26	104	25%	16%	37%		19	71	27%	16%	42%		22	89	25%	16%	37%		22	80	28%	17%	42%		18	88	21%	12%	32%
ever tried a whole cigarette		44	201	22%	16%	29%		37	148	25%	18%	34%		28	183	15%	10%	22%		not asked						not asked				
smoked cigarettes in past 30 days		25	203	12%	8%	18%		15	149	10%	6%	17%		15	184	8%	5%	13%		13	158	8%	4%	14%		9	181	5%	2%	9%
smoked bidis (flavored cigs) in past 30 days*		5	78	6%	2%	15%		3	33	9%	2%	26%		4	71	6%	2%	14%		6	74	8%	3%	17%		3	92	3%	1%	9%
chewed tobacco or used snuff in past 30 days		9	202	4%	2%	8%		6	149	4%	1%	9%		9	184	5%	2%	9%		10	157	6%	3%	12%		2	181	1%	0%	4%
smoked cigars/cigarillos/little cigars in past month		8	88	9%	4%	18%		3	33	9%	2%	26%		4	71	6%	2%	14%		7	73	10%	4%	20%		2	91	2%	0%	8%
smoked clove cigarettes in past 30 days		4	77	5%	1%	13%		1	34	3%	0%	15%		5	71	7%	2%	16%		not asked					not asked					
used a pain killer to get high in past 30 days		not asked						9	149	6%	3%	11%		7.01	91	8%	3%	16%		7	159	4%	2%	9%		7	178	4%	2%	8%
adults in neighborhood think it is very wrong/wrong for youth to:																														
use marijuana		97	110	88%	72%	100%		68	74	92%	71%	100%		84	96	88%	70%	100%		69	83	83%	65%	100%		82	90	91%	72%	100%
drink alcohol		93	110	85%	68%	100%		61	74	82%	63%	100%		79	96	82%	65%	100%		70	84	83%	65%	100%		81	89	91%	72%	100%
smoke		98	110	89%	72%	100%		62	73	85%	65%	100%		82	96	85%	68%	100%		67	84	80%	62%	100%		84	90	93%	75%	100%
very hard/sort of hard to get beer/wine/hard liquor		54	109	50%	37%	65%		43	74	58%	42%	78%		54	95	57%	43%	74%		53	83	64%	48%	84%		65	87	75%	58%	95%
very hard/sort of hard to get cigarettes		63	109	58%	44%	74%		94	143	66%	53%	80%		47	94	50%	37%	67%		48	82	59%	43%	78%		68	88	77%	60%	98%
very hard/sort of hard to get marijuana		72	109	66%	52%	83%		58	74	78%	60%	100%		60	93	65%	49%	83%		51	83	61%	46%	81%		66	89	74%	57%	94%
youth drinking alcohol would get caught by police		34	108	31%	22%	44%		28	72	39%	26%	56%		34	89	38%	26%	53%		27	84	32%	21%	47%		42	88	48%	34%	65%
youth smoking marijuana would get caught by police		52	106	49%	37%	64%		35	73	48%	33%	67%		43	91	47%	34%	64%		21	83	25%	16%	39%		45	88	51%	37%	69%
perceived great risk of harm from:																														
smoking a pack or more cigarettes a day		69	107	64%	50%	82%		51	74	69%	51%	91%		66	93	71%	55%	90%		60	84	71%	55%	92%		68	90	76%	59%	96%
smoking marijuana at least once/twice week		69	105	66%	51%	83%		51	74	69%	51%	91%		47	92	51%	38%	68%		43	84	51%	37%	69%		49	90	54%	40%	72%
drinking 1-2 alcoholic drinks daily		29	106	27%	18%	39%		20	74	27%	17%	42%		19	92	21%	12%	32%		27	84	32%	21%	47%		21	89	24%	15%	36%
perceived definite risk if youth:																														
smoke from 1-5 cigarettes per day		61	97	63%	48%	81%		52	75	69%	52%	91%		55	92	60%	45%	78%		54	74	73%	55%	95%		not asked				
breathe secondhand smoke		49	91	54%	40%	71%		45	72	63%	46%	84%		41	78	53%	38%	71%		36	64	56%	39%	78%		45	87	52%	38%	69%
school has staff to discuss substance use problem		79	107	74%	58%	92%		101	148	68%	56%	83%		96	172	56%	45%	68%		97	151	64%	52%	78%		104	178	58%	48%	71%
not sure school has staff to discuss sub use problem		22	107	21%	13%	31%		36	148	24%	17%	34%		57	172	33%	25%	43%		41	151	27%	20%	37%		63	178	35%	27%	45%

\*2010, 2012 question changed: "tobacco that tastes like candy, fruit or alcohol (little cigars, bidis, cloves, chew, spit, snus, hookah)"

### Response Rates:

70% or greater participation: results probably representative  
 40-69% participation: results may be representative  
 Less than 40% participation: results likely not representative

# Substance Use, 8th & 10th graders, Jefferson County: 2004, 2006, 2008, 2010, 2012

10th grade	2004			95% confidence interval		2006	2008			95% confidence interval		2010	2012			95% confidence interval									
	response rate: 61%	#	total	%	lower		upper	49%	#	total	%		lower	upper	53%	#	total	%	lower	upper					
ever had more than a sip or two of alcohol	116	178	65%	54%	78%	103	139	74%	61%	90%	96	138	70%	56%	85%	102	148	69%	56%	84%	71	109	65%	51%	82%
binge drank at least once in past 2 weeks	46	180	26%	19%	34%	17	64	27%	15%	42%	28	139	20%	13%	29%	31	148	21%	14%	30%	22	110	20%	13%	30%
drunk or high at school in past year	38	180	21%	15%	29%	41	144	28%	20%	39%	31	139	22%	15%	32%	40	148	27%	19%	37%	28	109	26%	17%	37%
drank alcohol in past 30 days	80	180	44%	35%	55%	65	147	44%	34%	56%	47	141	33%	24%	44%	55	150	37%	28%	48%	30	110	27%	18%	39%
ever smoked marijuana	81	181	45%	36%	56%	64	143	45%	34%	57%	51	138	37%	28%	49%	71	148	48%	38%	61%	53	109	49%	36%	64%
smoked marijuana in past 30 days	51	181	28%	21%	37%	44	147	30%	22%	40%	30	141	21%	14%	30%	51	150	34%	25%	45%	34	112	30%	21%	42%
ever tried a cigarette (even a puff)	38	83	46%	32%	63%	30	63	48%	32%	68%	30	68	44%	30%	63%	33	79	42%	29%	59%	19	55	35%	21%	54%
ever tried a whole cigarette	76	179	42%	33%	53%	54	141	38%	29%	50%	38	139	27%	19%	37%	not asked					19	54	35%	21%	55%
smoked cigarettes in past 30 days	41	181	23%	16%	31%	36	148	24%	17%	34%	19	140	14%	8%	21%	21	150	14%	9%	21%	17	112	15%	9%	24%
smoked bidis (flavored cigs) in past 30 days*	13	73	18%	9%	30%	2	28	7%	1%	25%	6	62	10%	4%	21%	7	69	10%	4%	21%	8	56	14%	6%	28%
chewed tobacco or used snuff in past 30 days	28	181	15%	10%	22%	13	148	9%	5%	15%	7	141	5%	2%	10%	8	150	5%	2%	10%	8	112	7%	3%	14%
smoked cigars/cigarillos/little cigars in past month	18	78	23%	14%	36%	5	28	18%	6%	41%	9	62	15%	7%	27%	10	69	15%	7%	27%	9	56	16%	7%	30%
smoked clove cigarettes in past 30 days	7	74	9%	4%	19%	5	28	18%	6%	41%	4	61	7%	2%	16%	not asked					not asked				
used a pain killer to get high in past 30 days	not asked					28	145	19%	13%	28%	9	72	13%	6%	24%	12	149	8%	4%	14%	11	109	10%	5%	18%
adults in neighborhood think it is very wrong/wrong for youth to:																									
use marijuana	70	88	80%	62%	100%	50	71	70%	52%	93%	56	76	74%	56%	96%	51	80	64%	48%	84%	40	56	71%	51%	97%
drink alcohol	57	88	65%	49%	84%	49	70	70%	52%	93%	48	76	63%	47%	84%	58	80	73%	55%	94%	43	56	78%	56%	104%
smoke	64	88	73%	56%	93%	55	72	76%	58%	99%	56	76	74%	56%	96%	60	80	75%	57%	97%	42	55	76%	55%	103%
very hard/sort of hard to get beer/wine/hard liquor	31	89	35%	24%	49%	17	72	24%	14%	38%	29	76	38%	26%	55%	27	81	33%	22%	48%	27	56	48%	32%	70%
very hard/sort of hard to get cigarettes	26	88	30%	19%	43%	32	115	28%	19%	39%	26	76	34%	22%	50%	28	81	35%	23%	50%	29	56	52%	35%	74%
very hard/sort of hard to get marijuana	38	87	44%	31%	60%	24	72	33%	21%	50%	34	76	45%	31%	63%	28	81	35%	23%	50%	20	56	36%	22%	55%
youth drinking alcohol would get caught by police	34	108	31%	22%	44%	28	72	39%	26%	56%	34	89	38%	26%	53%	19	81	23%	14%	37%	15	56	27%	15%	44%
youth smoking marijuana would get caught by police	20	87	23%	14%	35%	6	70	9%	3%	18%	13	74	18%	9%	30%	17	81	21%	12%	34%	16	56	29%	16%	46%
perceived great risk of harm from:																									
smoking a pack or more cigarettes a day	58	88	66%	50%	85%	41	70	59%	42%	80%	55	75	73%	55%	96%	53	81	65%	49%	86%	48	57	84%	62%	112%
smoking marijuana at least once/twice week	46	88	52%	38%	70%	27	69	39%	26%	57%	27	75	36%	24%	52%	18	81	22%	13%	35%	24	57	42%	27%	63%
drinking 1-2 alcoholic drinks daily	31	88	35%	24%	50%	18	68	26%	16%	42%	18	75	24%	14%	38%	21	81	26%	16%	40%	25	57	44%	28%	65%
perceived definite risk if youth:																									
smoke from 1-5 cigarettes per day	58	95	61%	46%	79%	55	80	69%	52%	90%	51	68	75%	56%	99%	51	69	74%	55%	97%	not asked				
breathe secondhand smoke	57	84	68%	51%	88%	28	51	55%	37%	79%	49	66	74%	55%	98%	43	69	62%	45%	84%	30	54	56%	38%	79%
school has staff to discuss substance use problem	65	88	74%	57%	94%	83	133	62%	50%	77%	77	139	55%	44%	69%	106	149	71%	58%	86%	78	109	72%	57%	89%
not sure school has staff to discuss substance use problem	15	88	17%	10%	28%	36	133	27%	19%	37%	48	139	35%	25%	46%	33	149	22%	15%	31%	23	109	21%	13%	32%

\*2010 and 2012 question changed: "tobacco that tastes like candy, fruit or alcohol (little cigars, bidis, cloves, chew, spit, snus, hookah)"

Data Sources: Washington State Department of Health, Healthy Youth Survey

Response Rates:

70% or greater participation: results probably representative

40-69% participation: results may be representative

Less than 40% participation: results likely not representative